

# *In-vivo* study of the anti-diabetic effect of *Ballota saxatilis*

Maha H. Abdulla <sup>1</sup>; Khlood A. Al-Khafaji <sup>2</sup>; Raghad S. Mouhamad <sup>2</sup>; Risala H. Allami <sup>1</sup>

1, College of Biotechnology, Al-Nahrain University, Baghdad, Iraq  
2, Ministry of Science and Technology, Baghdad, Iraq

## Abstract

**E-mail:**  
[raghad1974@yahoo.com](mailto:raghad1974@yahoo.com)

**Received:** 30/04/2020  
**Acceptance:** 07/06/2020  
**Available Online:** 07/06/2020  
**Published:** 01/07/2020

**Keywords:** *Ballota*, Fasting blood glucose; Diabetic Militias

Diabetes mellitus (DM) is recognized for years as a major public health problem with rapid expansion in developing countries. The main aim of the recent study is to evaluate the efficacy of the anti-diabetic effect of the aquatic extract of *Ballota saxatilis*. The hot aqueous extract was prepared from washed and air-dried aerial parts of the plant and the fatty acids in the extract were detected using gas chromatography system (GC). The extract was filtered and dried at 37°C to powder. Then, different concentrations were prepared using distilled water and orally injected into six groups of albino male mice for 21 days. Blood glucose was measured before and after treatment. The effects of *Ballota* extract (50 mg/Kg/day) on plasma glucose levels of normal mice was considerably more pronounced after four hours of application and the most significant decrease was observed after 21 days. A significant decrease in Fasting Serum Glucose (FSG) levels was observed under all *B. Saxatilis* extract applications in n5-Streptozotocin (n5-STZ) diabetic induced mice by the end of the experiment when compared to control. The most promising treatment was 150 mg/Kg/day as the subjects' FSG levels reached its normal level after 21 days of application. Therefore, due to its anti-hyperglycemic activity, the aerial parts extract of *ballota* can be considered a promising dietary supplement for type 2 diabetes patients especially those with a "pre-diabetic" state as a diabetes prevention method.

## 1. Introduction

Diabetes mellitus (DM) is a major health issue [1]. It is expected that the number of diabetic patients will increase to affect 7.7% of the population by 2030 [2]. Diabetes is associated with changes in the structure and functionality of the myocardium and this state is called diabetic cardiomyopathy (DCM) [3]. Severe complications are associated with diabetes such as cardiovascular disease and cerebrovascular disease, diabetic foot disorders, In addition to various microvascular disorders such as nephropathy, retinopathy, neuropathy, and sexual dysfunction [4].

Clinical and experimental diabetes mellitus results indicate that free radical oxidative stress mediates an important role in organ transformation and cardiac dysfunction [5]. Additionally, both type 1 and type 2 diabetes cause myocyte damage that occurs through reactive oxygen species (ROS) pathways [5][6].

While several plants have a proofed anti-diabetic effect, both research and conventional usage are limited to only a few useful plants, such as *Sylvestre gymnama*, neem, *Hordeum vulgare*, *Pterocarpus marsupium* and *Ocimum sanctum* [4]. *Ballota* also is known as Horehound (*Ballota saxatilis*) is a member of the family Lamiaceae and grows in various regions of Egypt's North Sinai [7]. This plant has a long history in traditional medicine, especially in Sinai for treating patients with non-insulin-dependent diabetes [8]. *Ballota* extract is rich in bioactive compounds such as methoxylated flavones, flavone aglycones, alkaloids, tannins, saponins, coumarins, docosanoic acid, catechins, amino acids, cardenolides, hydrocarbons, in addition to the sterols  $\beta$ -sitosterol and stigmasterol [9]. Moreover, the phytochemical screening of *ballota* has suggested the presence of 3- butenyl isothiocyanate, 2-methyl butenyl isothiocyanate, benzyl isothiocyanate, in addition to  $\alpha$ ,  $\beta$ , and 2-naphthyl-n-propyl ether in its volatile oil [10]. The phenolic compounds undoubtedly play a major role as antioxidants and free radical scavengers [11].



Therefore, the aim of this work is objected to investigate the probable hypoglycemic properties of ballota on diabetes mellitus in relation to its antioxidant effect in n5-Streptozotocin (n5-STZ)-induced diabetic mice.

## 2. Material and Methods

### 2.1. Plant material

The aerial part (stem) of Ballota was obtained from Jordan. Plant parts were washed thoroughly with distilled water to remove dirt and soil; then, they were air-dried for 72 hours at 25±2 °C.

### 2.2. Extract preparation

The aqueous extract was prepared by boiling 50g of ballota (stem)with 300 ml of distilled water for 5 minutes. After being cooled to room temperature, the mixture was placed in a shaker at room temperature for 24 h. The mixture was then filtered twice, first through a Whatman No. 0.4 filter paper and then through cotton wool. The aqueous extract was then poured in shallow layers in evaporating dishes and then incubated for 2-4 days at 37°C to obtain a powder extract of the plant. The powder extract was then diluted with distilled water to make different concentrations [12].

### 2.3. Gas chromatography-mass spectrometry (GC-MS) analysis

The fatty acids were detected in plant extracts using a gas chromatography system (GC)(Chrompack-Packard 438A) equipped with an FID Detector at a temperature of 325°C and a separation column type 30-SE with an inner diameter of 0.25 mm and a length of 30 m based on the method described in [13].

### 2.4. Laboratory Animals

The experiment was performed at the animal laboratory of the Biotechnology Research Center at Al-Nahrin University between March 2019 and October 2019. A total of 60 albino male mice weighing between 23-28 g were employed for this study. Mice were adapted for one week before the start of the experiment. The experimental design included twenty normal while the remained forty mice were given a single dose of of160mg/kg ofn5-Streptozotocin to induce neonatal diabetes [14]. The experimental animals were then divided into six different experimental groups, and each group was housed separately in transparent plastic cages equipped with stainless steel cover lids. The animals were maintained at a temperature of 25±2 °C, and they had ad libitum access to food (standard pellets) and water throughout the experimental period (21 days).

### 2.5. Experimental Design

The experiment was designed to evaluate the anti-diabetic effect of Ballota (water extract) in albino male mice. The tested groups were injected orally with a single dose (0.1ml)of different final concentrations (0, 50, 100, 150 mg/Kg) of Ballota extract per day for 21 days. The experimental groups are illustrated in (Table 1).

### 2.6. Blood sampling and determination of glucose level

Postprandial glucose level was monitored for treated normal mice (Group II) after 2 and 4 h of treatment on the first day, after one week, and at the end of the experiment. Fasting blood glucose levels of the experimental groups were monitored on the first day and at the end of the experiment. Blood samples were collected from the tail vein according to the method of [15] and blood samples were centrifuged for 10 minutes at 5000 rpm to separate the serum.

**Table 1. The different groups of mice used in the experiment with a final concentration of Ballota extract used in each group**

Group	Mice type	Treatment
Group I	Normal mice	Distilled water (Negative control)
Group II	Normal mice	50 mg/kg/day extract
Group III	Diabetic mice*	Distilled water (Positive control)
Group IV	Diabetic mice*	50 mg/kg/day extract
Group V	Diabetic mice*	100 mg/kg/day extract
Group VI	Diabetic mice*	150 mg/kg/day extract

\* Neonatal n5 Streptozotocin-induced DM mice

## 2.7. Measurement of glucose level

The glucose level in serum was determined immediately using a colorimetric method. First,  $\beta$ -D glucose is oxidized into gluconic acid and hydrogen peroxide in the presence of glucose oxidase (GOD).  $H_2O_2$  reacts with phenol and 4-amino antipyrine. The reaction forms a pink-colored quino amine dye complex. The glucose level is then estimated by measuring the absorbance of samples and standards against the blank (Distilled water) at 500 nm using a UV-VIS spectrophotometer (Unico –USA).

## 2.8. Statistical analysis

Data were analyzed with SAS Version 9.1 software using a one-way analysis of variance (ANOVA) with Tukey's HSD post hoc test at  $P < 0.05$ . All data results were presented as mean values ( $\pm$  SD) [16].

## 3. Results and Discussion:

The GC-MS analysis revealed that the aqueous hot extract of *Ballota* contained a different important type of saturated and unsaturated fatty acids. The highest concentration belonged to  $\alpha$ -Linolenic acid (Omega-3) followed by Palmitic acid with 57.05% and 27.28% respectively (Table 2). These results are in accordance with the results of [17].


**Table 2. fatty acids analysis of aquatic *Ballota* extract using GC/MS**

Fatty acid type	Name of fatty acid	The chemical formula of fatty acid		%
Saturated	Myristic acid	$C_{14}H_{28}O_2$	C14:0	1.22
	Palmitic acid	$C_{16}H_{32}O_2$	C16:0	27.28
Unsaturated	Linoleic acid (omega-6)	$C_{18}H_{32}O_2$	C18:3	3.15
	Oleic acid (omega-9 )	$C_{18}H_{34}O_2$	C18:1	2.48
	$\alpha$ -Linolenic acid (omega-3 )	$C_{18}H_{32}O_2$	C18:2	57.05
	Arachidonic acid	$C_{20}H_{32}O_2$	C20:4	2.56

The effects of *Ballota* extract (50 mg/Kg/day) on plasma glucose levels of normal mice after 2 and 4 hours of its oral application were investigated. Results illustrated that postprandial glucose level in normal mice (Group II) was reduced significantly by the end of the experiment after 21 days of application (Fig 1).

Numerous reports had suggested different species of the genus *Ballota* as a safe and inexpensive food supplement for various reasons [18][19]. The results of the current study illustrated the anti-hyperglycemic activity of *B. saxatilis* aquatic extract in normal mice which renders it a promising dietary supplement for those with "pre-diabetic" state as lifestyle activities including diet play a major role in DM development and control

Fasting serum glucose (FSG) levels were compared between the six mice models on the first day, and no differences were noticed between the different models except for those between normal (Groups I and II) and n5-STZ diabetic mice (Groups III, IV, V, and VI). After the oral application of *Ballota* extract for 21 days, the corresponding FSG level was significantly reduced in normal mice group with 50 mg/kg extract. A significant decrease in FSG levels was observed under all *B. Saxatilis* concentrations in n5-STZ diabetic mice (Group IV, Group V, and Group VI) by the end of the experiment when compared to control (Group III) (Fig. 2). The best treatment was 150 mg/Kg/day where the subjects FSG



**Figure 1. The effect of *Ballota* extract on postprandial glucose level (Mean $\pm$ SD) in normal mice (Group II)**


levels reached normal levels, with no significant difference between Group VI and Negative control (Group I) by the end of the experiment (Fig. 2).

The biological activity of *B. saxatilis* extract might be attributed to its unsaturated fatty acids content especially omega-3 which is known to improve several metabolic abnormalities linked to free radicals induced lipid peroxidation leading to diabetes mellitus as well as insulin-sensitizing effect leading to high production and secretion of adipocytokines as adiponectin and leptin. Additionally, omega-3 enhances fatty acid oxidation and reduces de novo lipogenesis which leads to a reduction in hepatic fat accumulation and preserves hepatic insulin sensitivity [20][21][22][23]. Another proposed theory to the anti-diabetic effect of omega-3 is attributed to the secretion of glucagon-like peptide-1 which is mediated by GPR120 [24]. Many studies reported that long-chain monounsaturated fatty acid, as well as  $\alpha$ -linolenic acid-containing diets, promote and stimulate GLP-1 secretion in murine rat and human L cells, and increase insulin circulating [25-29]. Additionally, [30] suggested foods of high n-3 PUFA content as a treatment strategy.

The aqueous extract of *Ballota saxatilis* appears to have promising antihyperglycemic effects in diabetes which might exceed the previous anti-diabetic activities in other *Ballota* genus members such as *B. nigra* [31][32] and *B. undulata* [33]. Therefore, the current study suggests *B. saxatilis* as a novel oral antihyperglycemic agent. Furthermore, extended studies are needed to investigate the cultivation of the plant in the Iraqi environment and to introduce its extracts to dietary programs.

## References

1. Rahimi-Madiseh M, Malekpour-Tehrani A, Bahmani M, Rafieian-Kopaei M. The research and development on the antioxidants in prevention of diabetic complications. *Asian Pac. J. Trop. Med.* 2016;9(9):825-31. DOI
2. Shaw JE, Sicree RA, Zimmet PZ. Global estimates of the prevalence of diabetes for 2010 and 2030. *Diabetes Res. Clin. Pract.* 2010;87(1):4-14. DOI
3. Liu Q, Wang S, Cai L. Diabetic cardiomyopathy and its mechanisms: role of oxidative stress and damage. *J Diabetes Invest.* 2014;5(6):623-34. DOI
4. Hariharan RS, Venkataraman S, Sunitha P, Rajalakshmi S, Samal KC, Routray BM, Jayakumar RV, Baiju K, Satyavati GV, Muthuswamy V, Gupta AK. Efficacy of vijayasar (*Pterocarpus marsupium*) in the treatment of newly diagnosed patients with type 2 diabetes mellitus: a flexible dose double-blind multicenter randomized controlled trial. *Diabetol Croat.* 2005;34(1):13-20.
5. Alhabib S, Aldrainly M, Alfarhan A. An evolving role of clinical pharmacists in managing diabetes: evidence from the literature. *Saudi Pharm J.* 2016;24(4):441-6. DOI
6. MacIsaac RJ, Jerums G, Ekinici EI. Cardio-renal protection with empagliflozin. *Ann. Transl. Med.* 2016;4(20):409. DOI



**Figure 2.** The effect of *Ballota* stems extract on fasting blood glucose level in normal and n5-STZ diabetic mice; \* The groups from left to right are the same groups mentioned in (Table 1) from Group I – VI.

7. Boulos L. Flora of Egypt; Azollaceae-Oxalidaceae, Vol 1. Cairo, Al-Hadara, 1999:170.
8. Shahwan MJ, Al-Qirim TM, Daradka H. Effects of *Ballota undulata* on blood biochemical parameters and insulin in albino rat. Asian Biomed. 2009;3:171-5.
9. Motaal AA, Ezzat SM, Haddad PS. Determination of bioactive markers in *Cleome droserifolia* using cell-based bioassays for antidiabetic activity and isolation of two novel active compounds. Phytomedicine. 2011;19(1):38-41. DOI
10. YILMAZ BS, Ergene B, ÇİTOĞLU GS. Determination of Total *Ortho*-Dihydroxycinnamic Acid Derivatives and Flavonoid Contents of *Ballota* Species Growing in Turkey. Turk J Pharm Sci. 2015;12(1):67-74.
11. Najafian M, Ebrahim-Habibi A, Yaghmaei P, Parivar K, Larijani B. Core structure of flavonoids precursor as an antihyperglycemic and antihyperlipidemic agent: an *in vivo* study in rats. Acta. Biochim. Pol. 2010; 57(4):553-60. DOI
12. Hameed SI, Al-Shahwany AW, Salih SJ. Evaluation of the Activity of Some Plants Extracts on Thyroid Gland Regulation in Female Albino Rats. Iraqi j. sci. 2020;61(2):254-65. DOI
13. Tipirdamaz R, Guvenc A. Seed fatty acids composition of *Ballota cristata*. Chem. Nat. Compd. 2004;40(3):291-2. DOI
14. Wu KK, Huan Y. Streptozotocin-induced diabetic models in mice and rats. Curr Protoc Pharmacol. 2008;40(1):5-47. DOI
15. Helal EG, Khattab AM, Abou Aouf N, Abdallah IZ. Hypoglycemic and Antioxidant Effects of *Cleome Droserifolia* (Samwah) in Alloxan-Induced Diabetic Rats. Egypt. J. Hosp. Med. 2015;58(1):39-47. DOI
16. SAS.Statistical Analysis System, User's Guide.Statistical Version 9.1. SAS.Inst. Inc. Cary.N.C. USA. 2012.
17. Rigano D, Marrelli M, Formisano C, Menichini F, Senatore F, Bruno M, Conforti F. Phytochemical profile of three *Ballota* species essential oils and evaluation of the effects on human cancer cells. Nat. Prod. Res. 2017;31(4):436-44. DOI
18. Morteza-Semnani K, Ghanbarimasir Z. A review on traditional uses, phytochemistry and pharmacological activities of the genus *Ballota*. J. Ethnopharmacol. 2019;233:197-217. DOI
19. Seidel V, Bailleul F, Tillequin F. Terpenoids and phenolics in the genus *Ballota* L.(*Lamiaceae*). Recent Res. Devel. Phytochem. 1999;3:27-39.
20. Patel DK, Kumar R, Laloo D, Hemalatha S. Diabetes mellitus: an overview on its pharmacological aspects and reported medicinal plants having antidiabetic activity. Asian Pac J Trop Biomed. 2012;2(5):411-20. DOI
21. Tanaka N, Zhang X, Sugiyama E, Kono H, Horiuchi A, Nakajima T, Kanbe H, Tanaka E, Gonzalez FJ, Aoyama T. Eicosapentaenoic acid improves hepatic steatosis independent of PPAR $\alpha$  activation through inhibition of SREBP-1 maturation in mice. Biochem pharmacol. 2010;80(10):1601-12. DOI
22. Neschen S, Morino K, Dong J, Wang-Fischer Y, Cline GW, Romanelli AJ, Rossbacher JC, Moore IK, Regittnig W, Munoz DS, Kim JH. n-3 Fatty acids preserve insulin sensitivity *in vivo* in a peroxisome proliferator-activated receptor- $\alpha$ -dependent manner. Diabetes. 2007;56(4):1034-41. DOI
23. Sato A, Kawano H, Notsu T, Ohta M, Nakakuki M, Mizuguchi K, Itoh M, Suganami T, Ogawa Y. Antiobesity effect of eicosapentaenoic acid in high-fat/high-sucrose diet-induced obesity: importance of hepatic lipogenesis. Diabetes. 2010;59(10):2495-504. DOI
24. Jump DB. Fatty acid regulation of hepatic lipid metabolism. Curr. Opin. Clin. Nutr. Metab. 2011 DOI
25. Lim GE, Brubaker PL. Glucagon-like peptide 1 secretion by the L-cell: the view from within. Diabetes. 2006;55(Supplement 2):S70-7. DOI

26. Reimer RA, Darimont C, Gremlich S, Nicolas-Métral V, Rüegg UT, Macé K. A human cellular model for studying the regulation of glucagon-like peptide-1 secretion. *Endocrinology*. 2001;142(10):4522-8. [DOI](#)
27. Allami RH, Mohsin RH, Mouhamad RS. Study the effect of herbal mixture plants extract on blood sugar level in normal and experimentally diabetic mice. *Revista Bionatura*. 2019;4(4):986-90. [DOI](#)
28. Mohsin RH, Allami RH, Mouhamad RS. X-ray fluorescence technique for studying mineral nutrients of Quinoa seed cultivated in Iraq. *Revista Bionatura*. 2019;4(4):966-71. [DOI](#)
29. Allami RH, Mouhamad RS, Mohsin RH. Determination of the physical, chemical and sensory properties of gluten-free biscuits by mixing different proportions of amber rice flour and quinoa flour. *Agri. Rech. J.* 2020;6(2): 20-5.
30. Hirasawa A, Tsumaya K, Awaji T, Katsuma S, Adachi T, Yamada M, Sugimoto Y, Miyazaki S, Tsujimoto G. Free fatty acids regulate gut incretin glucagon-like peptide-1 secretion through GPR120. *Nat. Med.* 2005;11(1):90-4. [DOI](#)
31. Nusier MK, Bataineh HN, Bataineh ZM, Daradka HM. Effects of *Ballota nigra* on glucose and insulin in alloxan-diabetic albino rats. *Neuro Endocrinol. Lett.* 2007;28(4):470-2.
32. Nusier MK, Bataineh HN, Bataineh ZM, Daradka HM. Effects of *Ballota nigra* on blood biochemical parameters and insulin in albino rats. *Neuro Endocrinol. Lett.* 2007;28(4):473-6.
33. Shahwan MJ, Al-Qirim TM, Daradka H. Effects of *Ballota undulata* on blood biochemical parameters and insulin in albino rat. *Asian Biomed.* 2009;3:171-5.